arXiv:0911.2408v2 [math.GR] 17 Nov 2009

Highly Transitive Actions of Surface Groups

Daniel Kitroser

November 17, 2009

Abstract

A group action is said to be highly-transitive if it is k-transitive for every $k \geq 1$. The main result of this thesis is the following:

Main Theorem. *The fundamental group of a closed, orientable surface of genus* > 1 *admits a faithfull, highly-transitive action on a countably infinite set.*

From a topological point of view, finding a faithfull, highly-transitive action of a surface group is equivalent to finding an embedding of the surface group into $\mathrm{Sym}(\mathbb{Z})$ with a dense image. In this topological setting, we use methods originally developed in [3] and [1] for densely embedding surface groups in locally compact groups.

1 Introduction

A permutation group $G \leq \mathrm{Sym}(X)$ is called *k-transitive* if it is transitive on ordered k-tuples of distinct elements and *highly-transitive* if it is k-transitive for every $k \in \mathbb{N}$. When G is given as an automorphism group, it can often be verified that the given action is highly transitive. This is the case for any subgroup of $\mathrm{Sym}(X)$ which contains all finitely supported permutations, where X is any infinite set. Other examples include groups such as $\mathrm{Homeo}(S^2)$ and many of its subgroups. But for most groups the question: "what is the maximal k for which the group admits a faithfull, k-transitive action"? is wide open.

It was shown by McDonough [4] that any non-abelian free group admits a highly transitive action. Another proof which is more useful from our point of view was later given by Dixon in [2], using a Baire Category type argument.

In this paper, we prove that a *surface group* (the fundamental group of a closed, orientable surface without boundry of genus greater than 1) admits a faithfull, highly-transitive action of countably infinite degree.

1.1 Topological interpretation

Before we continue, we introduce some notation.

Notation. We denote by $S_\infty = \mathrm{Sym}(\mathbb{Z})$ the full symmetric group of a countable set which we identify with the integer numbers. The action of $G \leq S_\infty$ on $\mathbb{Z}$ is always from the right and the image of an element $a \in \mathbb{Z}$ under $g \in G$ is denoted by a^g. If $n \in \mathbb{N}$ then $[n]$ will denote the set of all integers between $-n$ and n.

We define a group topology on S_∞ where a basis is given by the sets of the form

$$U(\varphi, [n]) = \left\{ \psi \in S_\infty \;\middle|\; \psi\big|_{[n]} = \varphi\big|_{[n]} \right\}, \ (\varphi \in S_\infty, n \in \mathbb{N}).$$

It is easy to see that if $\varphi \in S_\infty$ and $A \subseteq \mathbb{Z}$ is any finite set then $U(\varphi, A)$ is open in this topology. Notice that this topology is the restriction to S_∞ of the topology of pointwise convergence on the space of all functions $\mathbb{Z} \to \mathbb{Z}$ and a straightforward proof shows that S_∞ is a topological group which is completely metrizable.

The problem of finding highly-transitive subgroups of S_∞ can now be approached via the following proposition, derived directly from the definition of the topology on S_∞.

Proposition 1.1. *Let $G \leq S_\infty$. Then G is highly-transitive if and only if G is dense in S_∞.*

1.2 The main result

In view of proposition 1.1, the main result can be reformulated as follows.

Theorem 1.2 (main theorem)**.** *Let $\Gamma = \pi_1(\Sigma_g)$ be the fundamental group of an orientable surface of genus ≥ 2. Then there exists a dense subgroup of S_∞ which is isomorphic to Γ.*

The methods used in this work to obtain dense embeddings of surface groups into S_∞ are analogous to those used in [3] and [1] to show that if a locally compact group contains a dense, free subgroup of every finite rank > 1 then it contains a dense surface group of every genus > 1. S_∞ is *not* locally-compact but as we shall see, by prooving that S_∞ has a dense free subgroup of every finite rank > 1 with the additional property that certain elements of that free group generate a non-discrete cyclic group we can apply the same methods in our setting.

2 Dense Embeddings of Free Groups

As a first step to proving theorem 1.2 we prove the following.

Theorem 2.1. *Let* $r \geq 2$*, let* $w_i = w_i(\tau_1, \ldots, \tau_r)$*,* $(i \in \mathbb{N})$ *be reduced words and assume that there exists* $j \in \{1, \ldots, r\}$ *such that every* w_i *is not a conjugate of a power of* τ_j*. Then, there exist* $\tau_1, \ldots, \tau_r \in S_\infty$ *such that* $F = \langle \tau_1, \ldots, \tau_r \rangle$ *is a dense, rank r free subgroup of* S_∞ *and such that* $\langle w_i(\tau_1, \ldots, \tau_r) \rangle$ *is non-discrete for every* $i \in \mathbb{N}$*.*

We prove the theorem using Baire's Category Theorem.

Definition 2.2. Let X be a topological space. A subset of X which is a countable intersection of dense, open sets is called *residual* or *generic.*

Baire's theorem states that in a non-trivial, complete metric space, a residual set is dense. Since they are dense and closed under countable intersections, residual sets in a complete metric space can be thought of as being "large". Dixon showed in [2] that if we denote

$$U = \{(\tau_1, \ldots, \tau_r) \in S_\infty^r \mid \langle \tau_1, \ldots, \tau_r \rangle \text{ is dense in } S_\infty\}$$

then the set of elements in U that freely generate a free group is residual in $\overline{U}$. We use a somewhat different setting than Dixon in order to prove the existence of the desired dense free subgroups. Fix $\sigma \in S_\infty$ to be the *shift* permutation, i.e. $\forall a \in \mathbb{Z} : a^\sigma = a + 1$. We show that for every $n \geq 1$, the set of elements $(\tau_1, \ldots, \tau_n) \in S_\infty^n$ such that $\langle \sigma, \tau_1, \ldots, \tau_n \rangle$ has the properties stated in theorem 2.1 is residual is S_∞^n.

Lemma 2.3. *Let* $\gamma \in S_\infty$*.* $\langle \gamma \rangle \leq S_\infty$ *is non-discrete if and only if the orbits of* $\langle \gamma \rangle$ *are all finite and of unbounded length.*

Proof. Assume first that $\langle \gamma \rangle$ has an infinite orbit Δ and let $a \in \Delta$. Then, $\langle \gamma \rangle \cap U(1, \{a\}) = \{1\}$ and so $\langle \gamma \rangle$ is discrete. Now assume that the length of the orbits of $\langle \gamma \rangle$ is bounded. Let m be the product of the lengths of the orbits of $\langle \gamma \rangle$. So, $\gamma^m = 1$ thus, $\langle \gamma \rangle$ is finite and hence discrete.

Conversely, suppose that the orbits of $\langle \gamma \rangle$ finite and of unbounded length. To prove that $\langle \gamma \rangle$ is non-discrete it is enough to show that every basic neighborhood of the identity contains some non-trivial power of γ. Let $n \in \mathbb{N}$ and let $\Delta = \cup_{i=1}^k \Delta_i$ be a finite union of orbits of $\langle \gamma \rangle$ such that $[n] \subseteq \Delta$. By hypothesis, all the Δ_i's are finite. If we set $m = \prod_{i=1}^k |\Delta_i|$ then for every $a \in \Delta$ (and in particular for every $a \in [n]$) we have that $a^{\gamma^m} = a$ and since the orbit lengths of $\langle \gamma \rangle$ are unbounded, there is an orbit of $\langle \gamma \rangle$

which is longer then m so γ^m is not the identity element. Thus, $\gamma^m \neq 1$ is an element of the pointwise stablilizer $U(\mathrm{id}, [n])$. Since such stabilizers form a basis at the identity, we are finished. $\square$

Definition 2.4. Let $\gamma_1, \ldots, \gamma_n \in S_\infty$ and let $w = w(\gamma_1, \ldots, \gamma_n)$ be any word. If $w = w_1 w_2 \cdots w_n$ where $w_i \in \{\gamma_1^{\pm 1}, \ldots, \gamma_n^{\pm 1}\}$ then the *trace* of an element $a \in \mathbb{Z}$ under w is the ordered set

$$\mathrm{tr}_w(a) = \{a, a^{w_1}, a^{w_1 w_2}, \ldots, a^{w_1 w_2 \cdots w_n} = a^w\}.$$

Lemma 2.5. *Fix $n \in \mathbb{N}$ and let $w = w(\sigma, \tau_1, \ldots, \tau_n)$ be a reduced word which is not conjugated to a power of σ. Then the following sets are residual:*

(1) $\mathcal{F} = \{(\tau_1, \ldots, \tau_n) \in S_\infty^n \mid \langle \sigma, \tau_1, \ldots, \tau_n \rangle$ *is a free group of rank n* $\}$
(2) $\mathcal{D} = \{(\tau_1, \ldots, \tau_n) \in S_\infty^n \mid \langle \sigma, \tau_1, \ldots, \tau_n \rangle$ *is dense*$\}$
(3) $\mathcal{N} = \{(\tau_1, \ldots, \tau_n) \in S_\infty^n \mid \langle w(\sigma, \tau_1, \ldots, \tau_n,) \rangle$ *is non-discrete*$\}$

Proof. **(1)** Let v be a reduced, non-trivial word on $n+1$ letters and consider the set

$$\mathcal{F}_v = \{(\tau_1, \ldots, \tau_n) \in S_\infty^n \mid v(\sigma, \tau_1, \ldots, \tau_n) \neq 1\}.$$

If we prove that $\mathcal{F}_v$ is open and dense in S_∞^n for every v as above then $\mathcal{F}$ is residual since $\mathcal{F} = \bigcap_{v \neq 1} \mathcal{F}_v$. Obviously, $\mathcal{F}_v$ is open as the inverse image of the open set $S_\infty \smallsetminus \{1\}$ under the continuous mapping $(\tau_1, \ldots, \tau_n) \mapsto v(\sigma, \tau_1, \ldots, \tau_n)$. To prove that $\mathcal{F}_v$ is dense let $(\varphi_1, \ldots, \varphi_n) \in S_\infty^n$ and let $m \in \mathbb{N}$. We prove that there exists $(\tau_1, \ldots, \tau_n) \in \mathcal{F}_v$ such that $\tau_i\big|_{[m]} = \varphi_i\big|_{[m]}$ for every $1 \leq i \leq n$. First, write

$$v(\sigma, \tau_1, \ldots, \tau_n) = \sigma^{r_1} v_1 \sigma^{r_2} v_2 \cdots \sigma^{r_k} v_k \sigma^{r_{k+1}}$$

where $r_i \in \mathbb{Z}$ and $v_i \in \{\tau_1^{\pm 1}, \ldots, \tau_n^{\pm 1}\}$ for every i. We want define the permutations $\tau_1, \ldots, \tau_n$. Choose $a_1, \ldots, a_{k+1} \in \mathbb{Z}$ such that the numbers $a_1, \ldots, a_{k+1}, a_1 + r_1, \ldots, a_{k+1} + r_{k+1}$ are all distinct and define $(a_i + r_i)^{v_i} = a_{i+1}$ (e.g. if $v_1 = \tau_5^{-1}$ define $(a_1 + r_1)^{\tau_5^{-1}} = a_2$ or equivalently, define $a_2^{\tau_5} = a_1 + r_1$). Note that since all the integers involved in the definition of the v_i's are distinct and v is reduced, the definitions of the v_i's do not contradict each other. We now have that $a_1^v = a_{k+1} + r_{k+1}$ and in particular, v is not trivial. In order to fulfill the condition that τ_i must send an element $b \in [m]$ to b^{φ_i} we choose $a_1, \ldots, a_{k+1}$ in a way that will not contradict with this requirment. Explicitly, choose

$$a_1, \ldots, a_{k+1} \in \mathbb{Z} \smallsetminus \left([m] \cup \left(\bigcup_i [m]^{\varphi_i} \right) \cup \left(\bigcup_j [m] - r_j \right) \cup \left(\bigcup_{i,j} [m]^{\varphi_i} - r_j \right) \right).$$

Note that we can always choose $a_1, \dots, a_{k+1}$ in the manner described since we are only excluding a finite set of integers that we can not choose from. Now, every τ_i is defined on $[m]$ and on some other elements $\{b_1, \dots, b_\ell\} \subseteq \{a_1, \dots, a_k, a_1 + r_1, \dots, a_{k+1} + r_{k+1}\}$, sending them to $\{c_1, \dots, c_\ell\}$ respectively. Finally, choose any bijection

$$f_i : \mathbb{Z} \smallsetminus \big([m] \cup \{b_1, \dots, b_\ell\}\big) \to \mathbb{Z} \smallsetminus \big([m]^{\varphi_i} \cup \{c_1, \dots, c_\ell\}\big)$$

and define

$$x^{\tau_i} = \begin{cases} x^{\varphi_i} & , x \in [m]. \\ c_j & , x = b_j \text{ for some } j. \\ x^{f_i} & , \text{else}. \end{cases}$$

So we have that every τ_i is a permutation which lies in the basic neighbourhood of φ_i defined by $[m]$ such that $v(\sigma, \tau_1, \dots, \tau_n)$ is non-trivial.

(2) Fix some $k \in \mathbb{N}$ and for every two k-tuples $\mathbf{x} = (x_1, \dots, x_k)$ and $\mathbf{y} = (y_1, \dots, y_k)$ of distinct integers consider the set

$$\mathcal{D}_{\mathbf{x},\mathbf{y}} = \left\{ (\tau_1, \dots, \tau_n) \in S_\infty^n \;\middle|\; \begin{array}{l} \exists \varphi \in \langle \sigma, \tau_1, \dots, \tau_n \rangle : \\ \forall 1 \le j \le k : x_j^{\varphi} = y_j \end{array} \right\}.$$

Notice that if $(\tau_1, \dots, \tau_n) \in \bigcap_{\mathbf{x},\mathbf{y}} \mathcal{D}_{\mathbf{x},\mathbf{y}}$ where $\mathbf{x}$ and $\mathbf{y}$ range over all k-tuples of distinct integers then $\langle \sigma, \tau_1, \dots, \tau_n \rangle$ is k-transitive. Thus, by proposition 1.1 we have that

$$\mathcal{D} = \bigcap_{k \in \mathbb{N}} \bigcap_{\mathbf{x},\mathbf{y}} \mathcal{D}_{\mathbf{x},\mathbf{y}}$$

and we are left to show that $\mathcal{D}_{\mathbf{x},\mathbf{y}}$ is open and dense for every $\mathbf{x}$ and $\mathbf{y}$ as above.

First, fix $\mathbf{x} = (x_1, \dots, x_k), \mathbf{y} = (y_1, \dots, y_k)$ as above and let $(\tau_1, \dots, \tau_n) \in \mathcal{D}_{\mathbf{x},\mathbf{y}}$. By definition there exists $\varphi \in \langle \sigma, \tau_1, \dots, \tau_n \rangle$ such that $x_i^{\varphi} = y_i$ for every $i = 1, \dots, k$. Let us write $\varphi = v(\sigma, \tau_1, \dots, \tau_n)$ where $v = v(\sigma, \tau_1, \dots, \tau_n)$ is a word on $\{\sigma^{\pm 1}, \tau_1^{\pm 1}, \dots, \tau_n^{\pm 1}\}$ and define $A = \bigcup_{i=1}^{k} \operatorname{tr}_v(x_i)$. A is finite and so the set

$$\mathcal{U} = \left\{ (\psi_1, \dots, \psi_n) \in S_\infty^n \;\middle|\; \begin{array}{l} \psi_i\big|_A = \tau_i\big|_A,\ \psi_i^{-1}\big|_A = \tau_i^{-1}\big|_A \\ i = 1, \dots, n \end{array} \right\}$$

is an open neighbourhood of $(\tau_1, \dots, \tau_n)$ contained in $\mathcal{D}_{\mathbf{x},\mathbf{y}}$. Indeed, if $(\psi_1, \dots, \psi_n) \in \mathcal{U}$ take $\xi = v(\sigma, \psi_1, \dots, \psi_n) \in \langle \sigma, \psi_1, \dots, \psi_n \rangle$, then by the definition of $\mathcal{U}$, we have that ξ acts the same as $\varphi = v(\sigma, \tau_1, \dots, \tau_n)$ on x_i and in particular, ξ sends x_i to y_i. This shows that $\mathcal{D}_{\mathbf{x},\mathbf{y}}$ is open.

Now we prove that $\mathcal{D}_{\mathbf{x},\mathbf{y}}$ is dense. Let $(\varphi_1, \dots, \varphi_n) \in S_\infty^n$ and $m \in \mathbb{N}$. Let $r \in \mathbb{N}$ be such that $x_j + r \notin [m]$ and $y_j + r \notin [m]^{\varphi_1}$ for every $1 \leq j \leq k$. Let

$$f : \mathbb{Z} \setminus \big([m] \cup \{x_1 + r, \dots, x_k + r\}\big) \to \mathbb{Z} \setminus \big([m]^{\varphi_1} \cup \{y_1 + r, \dots, y_k + r\}\big)$$

be any bijection and define

$$x^{\tau_1} = \begin{cases} x^{\varphi_1} & , x \in [m] \\ y_j + r & , x = x_j + r \text{ for some } 1 \leq j \leq k \\ x^f & , \text{otherwise} \end{cases}.$$

Now define $\tau_i = \varphi_i$ for every $2 \leq i \leq n$ and we get that $(\tau_1, \dots, \tau_n)$ is an element of the basic neighbourhood of $(\varphi_1, \dots, \varphi_n)$ defined by $[m]$. Also, the permutation $\xi = \sigma^r \tau_1 \sigma^{-r} \in \langle \sigma, \tau_1, \dots, \tau_n \rangle$ sends each x_j to y_j thus, $(\tau_1, \dots, \tau_n) \in \mathcal{D}_{\mathbf{x},\mathbf{y}}$.

(3) By lemma 2.3 we can equivalently write

$$\mathcal{N} = \left\{ (\tau_1, \dots, \tau_n) \in S_\infty^n \;\middle|\; \begin{array}{l} \text{The orbits of } \langle w(\sigma, \tau_1, \dots, \tau_n) \rangle \text{ are all finite} \\ \text{and of unbounded length.} \end{array} \right\}.$$

Thus, if we define for every $t \in \mathbb{N}$ and $a \in \mathbb{Z}$:

$$\mathcal{U}_t = \{(\tau_1, \dots, \tau_n) \in S_\infty^n \mid \langle w(\sigma, \tau_1, \dots, \tau_n) \rangle \text{ has an orbit of length} \geq t\}.$$
$$\mathcal{V}_a = \{(\tau_1, \dots, \tau_n) \in S_\infty^n \mid \text{The orbit of } a \text{ under } \langle w(\sigma, \tau_1, \dots, \tau_n) \rangle \text{ is finite}\}.$$

we get that $\mathcal{N} = \big(\bigcap_{t \in \mathbb{N}} \mathcal{U}_t\big) \cap \big(\bigcap_{a \in \mathbb{Z}} \mathcal{V}_a\big)$.

We now show that $\mathcal{U}_t$ and $\mathcal{V}_a$ are open and dense for every $t \in \mathbb{N}$ and $a \in \mathbb{Z}$. Let $(\tau_1, \dots, \tau_n) \in \mathcal{U}_t$ and let $b \in \mathbb{Z}$ be an element belonging to an orbit of $\langle w(\sigma, \tau_1, \dots, \tau_n) \rangle$ of length $\geq t$. Thus, $b, b^w, b^{w^2}, \dots, b^{w^{t-1}}$ are all distinct. Let $\Delta = \bigcup_{i=1}^{k-1} \mathrm{tr}_{w^i}(b)$. The set

$$\left\{ (\psi_1, \dots, \psi_n) \in S_\infty^n \;\middle|\; \begin{array}{l} \psi_i\big|_\Delta = \tau_i\big|_\Delta, \ \psi_i^{-1}\big|_\Delta = \tau_i^{-1}\big|_\Delta \\ i = 1, \dots, n \end{array} \right\}$$

is an open neighbourhood of $(\tau_1, \dots, \tau_n)$ which is contained in $\mathcal{U}_t$ hence, $\mathcal{U}_t$ is open.

Now, take $(\tau_1, \dots, \tau_n) \in \mathcal{V}_a$ and let $\Delta = \{a_1, \dots, a_s\}$ be the finite orbit of $\langle w(\sigma, \tau_1, \dots, \tau_n) \rangle$ containing a. Similarly

$$\left\{ (\psi_1, \dots, \psi_n) \in S_\infty^n \;\middle|\; \begin{array}{l} \psi_i\big|_\Delta = \tau_i\big|_\Delta, \ \psi_i^{-1}\big|_\Delta = \tau_i^{-1}\big|_\Delta \\ i = 1, \dots, n \end{array} \right\}$$

is an open neighbourhood of $(\tau_1, \ldots, \tau_n)$ which is contained in $\mathcal{V}_a$.

To prove $\mathcal{U}_t$ is dense, let $(\varphi_1, \ldots, \varphi_n) \in S_\infty^n$ and $m \in \mathbb{Z}$. In (1) we in fact showed that we can define $\tau_1, \ldots, \tau_n \in S_\infty$ such that for every finite set $A \subseteq \mathbb{Z}$ we have that $\tau_i\big|_A$ acts in any way we please and there exists some $b \in \mathbb{Z}$ such that $b^w \neq b$ where $w = w(\sigma, \tau_1, \ldots, \tau_n)$. By repeating the same argument we can find $\tau_1, \ldots, \tau_n \in S_\infty$ such that $\tau_i\big|_{[m]} = \varphi_i\big|_{[m]}$ for every $1 \leq i \leq n$ and such that there exists $b \in \mathbb{Z}$ such that $b, b^w, b^{w^2}, \ldots, b^{w^{t-1}}$ are all distinct i.e. $\langle w(\sigma, \tau_1, \ldots, \tau_n)\rangle$ has an orbit of length $\geq t$.

Finally, we prove that $\mathcal{V}_a$ is dense. Since $\langle w(\sigma, \tau_1, \ldots, \tau_n)\rangle$ has the same orbit structure as the cyclic group generated by any conjugate of w, we can assume without loss of generality that w is a cyclically reduced word that is not a power of σ. Let $(\varphi_1, \ldots, \varphi_n) \in S_\infty^n$ and $m \in \mathbb{Z}$. We need to define permutations $\tau_1, \ldots, \tau_n \in S_\infty$ such that $(\tau_1, \ldots, \tau_n) \in \mathcal{V}_a$ and every τ_i agrees with φ_i on $[m]$. This condition can be thought of in the following way: τ_i is already defined on $[m]$ for every i and τ_i^{-1} is already defined on $[m]^{\varphi_i}$ for every i (they act the same as φ_i and φ_i^{-1} respectively) and we are left to define τ_i on $\mathbb{Z} \smallsetminus [m]$ (and τ_i^{-1} on $\mathbb{Z} \smallsetminus [m]^{\varphi_i}$) in such a way that the orbit of a under $\langle w(\sigma, \tau_1, \ldots, \tau_n)\rangle$ will be finite. First we write $w(\sigma, \tau_1, \ldots, \tau_n) = w_1 w_2 \cdots w_k$. Now, we start by applying the positive and negative powers of w to a letter by letter:

$$\xrightarrow{w_\ell} c \xrightarrow{w_{\ell+1}} \ldots \xrightarrow{w_{n-1}} a_{-1} \xrightarrow{w_n} a = a_0 \xrightarrow{w_1} a_1 \xrightarrow{w_2} \ldots \xrightarrow{w_{s-1}} b \xrightarrow{w_s}$$

where $b \in \mathbb{Z}$ is the first element such that we need to apply to it the permutation w_s and w_s is not yet defined on b, that is, $w_s = \tau_i$ for some i and $b \notin [m]$ or $w_s = \tau_i^{-1}$ for some i and $b \notin [m]^{\varphi_i}$. Note that if such an element b does not exist then, since by hypothesis at least one of the letters $w_1, \ldots, w_k$ is not σ, the orbit of a under $\langle w(\sigma, \varphi_1, \ldots, \varphi_n)\rangle$ is contained in $[m] \cup \left(\bigcup_{i=1}^n [m]^{\varphi_i}\right)$, hence finite and we can just take $\tau_i = \varphi_i$ for every i. We can thus assume that such a b exists. Similarly, $c \in \mathbb{Z}$ is the first element we reach when applying letter by letter the negative powers of w to a such that we need to apply to c the permutation w_ℓ^{-1} and w_ℓ^{-1} is not yet defined on c. As before, we can assume without loss of generality that such an element c exists. By hypothesis, w is cyclically reduced and so the word $w_s w_{s+1} \cdots w_k w_1 \cdots w_\ell$ is reduced. Also, by their definition, $w_s, w_\ell \neq \sigma^{\pm 1}$. We wish to define $\tau_1, \ldots, \tau_n$ in such a way that $b^{w_s w_{s+1} \cdots w_k w_1 \cdots w_\ell} = c$ (and of course fulfilling the condition that τ_i agrees with φ_i on $[m]$). By repeating the argument made in (1), we can find two distinct elements $d_1, d_2 \in \mathbb{Z}$ that do not lie in $[m]$ or any $[m]^{\varphi_i}$ such that $w_{s+1} \cdots w_k w_1 \cdots w_{\ell-1}$ sends d_1 to d_2 and define $b^{w_s} = d_1$, $d_2^{w_\ell} = c$. From this we get that $b^{w_s \cdots w_k w_1 \cdots w_l} = d_1^{w_{s+1} \cdots w_k w_1 \cdots w_\ell} = d_2^{w_\ell} = c$ thus, the orbit of a is finite. Now, every τ_i is defined on $[m]$ and maybe on finitely many more elements. Again, exactly as we did in (1), we can extend the definition of τ_i to $\mathbb{Z}$ and get permutations $\tau_1, \dots, \tau_n$ satisfying the required conditions. □

proof of theorem 2.1. Let $j \in \{1, \dots, r\}$ be such that every word w_i is not a conjugate of a power of τ_j. If we set $\tau_j = \sigma$ then by lemma 2.5, the set

$$W_i = \left\{ (\tau_1, \dots, \tau_{j-1}, \tau_{j+1}, \dots, \tau_r) \in S_\infty^{r-1} \;\middle|\; \begin{array}{l} \langle \tau_1, \dots, \tau_r \rangle \text{ is a dense, rank } r \text{ free subgroup} \\ \text{and } \langle w_i(\tau_1, \dots, \tau_r) \rangle \text{ is non-discrete} \end{array} \right\}$$

is residual and so $\bigcap_{i \in \mathbb{N}} W_i$ is residual and in particular, not empty. □

3 Eventually Faithfull Sequences

Definition 3.1. Let G and H be groups. A sequence $\{f_n\}_{n=1}^{\infty}$ of homomorphisms from G to H is *eventually faithfull* if for every $g \in G$ there exists $n_0 \in \mathbb{N}$ such that $g \notin \ker(f_n)$ for all $n \geq n_0$.

In order to prove the main theorem, we will need to produce eventually faithfull sequences of homomorphisms from surface groups to free groups. The following constructions apear in [3] and [1]. Let $\Gamma = \Gamma_{2r}$ be the surface group of genus $2r$ $(r \geq 1)$. We have the following presentation for Γ

$$\Gamma = \langle a_1, a_1' \dots, a_r, a_r', b_1, b_1', \dots, b_r, b_r' \mid [a_1, a_1'] \cdots [a_r, a_r'][b_r', b_r] \cdots [b_1', b_1] \rangle.$$

Let $x = [a_1, a_1'] \cdots [a_r, a_r']$ and let $h : \Gamma \to \Gamma$ be the Dehn twist around x, i.e.

$$\begin{aligned} h(a_i) &= a_i \qquad & h(b_i) &= x b_i x^{-1} \\ h(a_i') &= a_i' \qquad & h(b_i') &= x b_i' x^{-1} \end{aligned}$$

Let F be the free group on $2r$ free generators $\{\varphi_1, \varphi_1', \dots, \varphi_r, \varphi_r'\}$ and let $k : \Gamma \to F$ be the homomorphism defined by

$$\begin{aligned} k(a_i) = k(b_i) &= \varphi_i \\ k(a_i') = k(b_i') &= \varphi_i'. \end{aligned}$$

Consider the map that folds the genus $2r$ surface that has Γ as its fundamental group across the curve corresponding to x (this curve seperates the surface into two equal parts). The image of this map is a surface of genus r with one boundry component so it has F as its fundamental group. k is the homomorphism induced on the fundamental groups by this folding map (see figure 1). Denote $f_n = k \circ h^n$.

Lemma 3.2 (Breuillard,Gelander,Souto,Storm [3]). *The sequence* $\{f_n\}_{n=1}^{\infty}$ *is eventually faithfull.*

We now construct an eventually faithfull sequence of homomorphisms from an odd genus surface group into a free group. Let $\Gamma = \Gamma_{2r+1}$ be a surface group of genus $2r+1$ ($r \geq 1$) with the presentation

$$\Gamma = \langle a_1, a_1', \ldots, a_r, a_r', b, b', c_1, c_1', \ldots, c_r, c_r' \mid [a_1, a_1'] \cdots [a_r, a_r'][b', b][c_1', c_1] \cdots [c_r', c_r] \rangle$$

Denote $x = [a_1, a_1'] \cdots [a_r, a_r'] b'$ and let F be the free group on $2r+1$ free generators $\{\varphi_1, \varphi_1', \ldots, \varphi_r, \varphi_r', \tau\}$. Let $\delta : \Gamma \to \Gamma$ and $\zeta : \Gamma \to \Gamma$ be denh



Figure 1:

twists around x and b' respectively, that is

$$\begin{aligned}
\delta(a_i) &= a_i & \zeta(a_i) &= a_i \\
\delta(a_i') &= a_i' & \zeta(a_i') &= a_i' \\
\delta(b) &= xb & \zeta(b) &= b(b')^{-1} \\
\delta(b') &= b' & \zeta(b') &= b' \\
\delta(c_i) &= xc_ix^{-1} & \zeta(c_i) &= c_i \\
\delta(c_i') &= xc_i'x^{-1} & \zeta(c_i') &= c_i'
\end{aligned}$$

Notice that δ and ζ commute.



Figure 2:

Let $k : \Gamma \to F$ be the map induced by folding the $2r+1$ surface across the curves x and b' (these curves seperate the surface into two surfaces of genus r and two boundry components). Explicitely,

$$\begin{aligned} k(a_i) &= \varphi_i \\ k(a'_i) &= \varphi'_i \\ k(b) &= 1 \\ k(b'_i) &= \tau \\ k(c_i) &= \varphi_i \\ k(c'_i) &= \varphi'_i \end{aligned}$$

(see figure 2). Finally, denote $\rho_n = k \circ (\delta \circ \zeta)^n$.

Lemma 3.3 (Barlev,Gelander [1]). *The sequence* $\{\rho_n\}_{n=1}^{\infty}$ *is eventually faithfull.*

4 Proof of The Main Theorem

We will need the following results.

Lemma 4.1. *Let* $\varphi, \psi \in S_\infty$ *be such that both* $\langle \varphi \rangle$ *and* $\langle \psi \rangle$ *are non-discrete. Then* $\langle (\varphi, \psi) \rangle \leq S_\infty^2$ *is non-discrete.*

Proof. We prove that every basic neighbourhood of $(1,1)$ contains a non-trivial element of $\langle (\varphi, \psi) \rangle$. Let $m \in \mathbb{N}$. Let $\Delta = \bigcup_{i=1}^{\ell} \Delta_i$ and $\Gamma = \bigcup_{i=1}^{k} \Gamma_i$ be finite unions of orbits of $\langle \varphi \rangle$ and $\langle \psi \rangle$ respectively such that $[m] \subseteq \Delta$ and $[m] \subseteq \Gamma$. From proposition 2.3 we have that all the orbits of $\langle \varphi \rangle$ and $\langle \psi \rangle$ are finite and so we can define the number

$$n = \prod_{i=1}^{\ell} |\Delta_i| \cdot \prod_{i=1}^{k} |\Gamma_i|.$$

Notice that every element of Δ is fixed by φ^n and every element of Γ is fixed by ψ^n and in particular every $i \in [m]$ is fixed by φ^n and ψ^n. From proposition 2.3 we also have that the lengths of the orbits of $\langle \varphi \rangle$ and $\langle \psi \rangle$ are unbounded and in particular $\langle \varphi \rangle$ and $\langle \psi \rangle$ both have an orbit of length greater then n and so φ^n and ψ^n are non-trivial. Thus $(\varphi, \psi)^n$ is a non-trivial element contained in the basic neighbourhood of $(1,1)$ defined by $[m] \times [m]$. $\square$

Lemma 4.2. *Let G be a Hausdorff topological group and let $\gamma \in G$ such that $\langle \gamma \rangle$ is non-discrete. Then, for every $n_0 \in \mathbb{N}$ the set $\{\gamma^n \mid n \geq n_0\}$ is dense in $\overline{\langle \gamma \rangle}$.*

Proof. First we notice that since $\langle\gamma\rangle$ is non discrete then also $\overline{\langle\gamma\rangle}$ is non discrete. Let $U \subseteq \overline{\langle\gamma\rangle}$ be open then we have $\gamma^m \in U$ for some $m \in \mathbb{Z}$. If $m \geq n_0$ we are done so assume $m < n_0$. Now, If we denote

$$\begin{aligned} U' &= U\gamma^{-m} \\ U'' &= U' \cap (U')^{-1} \end{aligned}$$

Then U'' is an open symmetric identity neighborhood and since $\overline{\langle\gamma\rangle}$ is non discrete, U'' is not finite. Now take

$$\widetilde{U} = U'' \smallsetminus \{\gamma^k \ : \ |k| < n_0 - m\}.$$

We have that $\widetilde{U}$ is open (since G is Hausdorff) and non empty, thus there exits $n \in \mathbb{Z}$ such that $\gamma^n \in \widetilde{U}$. By the definition of $\widetilde{U}$ we have that $|n| \geq n_0 - m$. Also, since $\widetilde{U} \subseteq U''$ and U'' is symmetric it follows that $\gamma^n, \gamma^{-n} \in U'' \subseteq U'$. Hence, $\gamma^{n+m}, \gamma^{-n+m} \in U$ and since either $n + m \geq n_0$ or $-n + m \geq n_0$ we are done. $\square$

4.1 Proof of theorem 1.2 for even genus

By theorem 2.1 there exists a subgroup $F \leq S_\infty$ such that F is dense, free with $2r$ free generators $\varphi_1, \varphi'_1, \ldots, \varphi_r, \varphi'_r \in S_\infty$ and such that $\overline{\langle[\varphi_1, \varphi'_1] \cdots [\varphi_r, \varphi'_r]\rangle}$ is non-discrete. Denote $\gamma = [\varphi_1, \varphi'_1] \cdots [\varphi_r, \varphi'_r]$ and $\Omega = \overline{\langle\gamma\rangle}$. Let $\Gamma = \Gamma_{2r}$ be a surface group of genus $2r$ ($r \geq 1$) with the presentation

$$\Gamma = \langle a_1, a'_1 \ldots, a_r, a'_r, b_1, b'_1, \ldots, b_r, b'_r \mid [a_1, a'_1] \cdots [a_r, a'_r][b'_r, b_r] \cdots [b'_1, b_1]\rangle.$$

Define for every $\omega \in \Omega$ a homomorphism $f_\omega : \Gamma \to S_\infty$ by

$$\begin{aligned} f_\omega(a_i) &= \varphi_i \\ f_\omega(a'_i) &= \varphi'_i \\ f_\omega(b_i) &= \omega\varphi_i\omega^{-1} \\ f_\omega(b'_i) &= \omega\varphi'_i\omega^{-1}. \end{aligned}$$

Since every $\omega \in \Omega$ commutes with γ this defines a homomorphism. Indeed, for every $\omega \in \Omega$:

$$f_\omega([a_1, a'_1] \cdots [a_r, a'_r][b'_r, b_r] \cdots [b'_1, b_1]) = \underbrace{[\varphi_1, \varphi'_1] \cdots [\varphi_r, \varphi'_r]}_{\gamma}\, \omega \, \underbrace{[\varphi'_r, \varphi_r] \cdots [\varphi'_1, \varphi_1]}_{\gamma^{-1}}\, \omega^{-1} =$$

$$\gamma\omega\gamma^{-1}\omega^{-1} = 1.$$

Ω is a completely metrizable space and every element of Ω corresponds to a homomorphism $\Gamma \to S_\infty$ whose image contains F, hence the image is dense. We are left to show that at least one of those homorphisms is also faithfull. In fact, we show that $\chi = \{\omega \in \Omega \mid f_\omega \text{ is faithfull}\}$ is residual in Ω.

For every $g \in \Gamma \smallsetminus \{1\}$ denote $\chi_g = \{\omega \in \Omega \mid f_\omega(g) \neq 1\}$. Notice that

$$\chi = \bigcap_{g \in \Gamma \smallsetminus \{1\}} \chi_g.$$

From lemma 3.2, the sequence $\{f_{\gamma^n}\}_{n \in \mathbb{N}}$ is eventually faithfull and so for every $g \in \Gamma \smallsetminus \{1\}$ there exists $n_0 \in \mathbb{N}$ such that $\gamma^n \in \chi_g$ for all $n \geq n_0$. By lemma 4.2 we have that $\{\gamma^n \mid n \geq n_0\}$ is dense in Ω and so, χ_g is dense in Ω. χ_g is also open as the inverse image of the the open set $S_\infty \smallsetminus \{1\}$ under the continuous map $\omega \to f_\omega(g)$. This shows that χ is residual in Ω. $\square$

4.2 Proof of theorem 1.2 for odd genus

Let $\Gamma = \Gamma_{2r+1}$ be a surface group of genus $2r+1$ $(r \geq 1)$ with the presentation

$$\Gamma = \langle a_1, a_1', \ldots, a_r, a_r', b, b', c_1, c_1', \ldots, c_r, c_r' \mid [a_1, a_1'] \cdots [a_r, a_r'][b', b][c_1', c_1] \cdots [c_r', c_r] \rangle$$

Let $F \leq S_\infty$ be a dense, free subgroup with $2r+1$ free generators $\varphi_1, \varphi_1', \ldots, \varphi_r, \varphi_r', \tau \in S_\infty$ such that $\langle \tau \rangle$ and $\langle [\varphi_1, \varphi_1'] \cdots [\varphi_r, \varphi_r'] \tau \rangle$ are non-discrete (the existence of such a free subgroup is assured by theorem 2.1). Denote $\gamma = [\varphi_1, \varphi_1'] \cdots [\varphi_r, \varphi_r'] \tau$ and $\Omega = \overline{\langle (\gamma, \tau) \rangle} \subseteq S_\infty^2$. For every $(\psi, \xi) \in \Omega$ we define a homomorphism $f_{(\psi,\xi)} : \Gamma \to S_\infty$ by setting

$$\begin{array}{lll} f_{(\psi,\xi)}(a_i) = \varphi_i & f_{(\psi,\xi)}(b) = \psi\xi^{-1} & f_{(\psi,\xi)}(c_i) = \psi\varphi_i\psi^{-1} \\ f_{(\psi,\xi)}(a_i') = \varphi_i' & f_{(\psi,\xi)}(b') = \tau & f_{(\psi,\xi)}(c_i') = \psi\varphi_i'\psi^{-1} \end{array}$$

Since every $(\psi, \xi) \in \Omega$ commutes with (γ, τ) we have that $f_{(\psi,\xi)}$ is well defined as a homomorphism because

$$\begin{aligned} & f_{(\psi,\xi)}([a_1, a_1'] \cdots [a_r, a_r'][b', b][c_1', c_1] \cdots [c_r', c_r]) = \\ & = [\varphi_1, \varphi_1'] \cdots [\varphi_r, \varphi_r'][\tau, \psi\xi^{-1}]\psi \underbrace{[\varphi_1', \varphi_1] \cdots [\varphi_r', \varphi_r]}_{=\tau\gamma^{-1}} \psi^{-1} = \\ & = \gamma\psi\xi^{-1}\tau^{-1}\xi\psi^{-1}\psi\tau\gamma^{-1}\psi^{-1} = \gamma\psi\xi^{-1}\tau^{-1}\xi\tau\gamma^{-1}\psi^{-1} = \\ & = \gamma\psi\xi^{-1}\xi\tau^{-1}\tau\gamma^{-1}\psi^{-1} = \gamma\psi\gamma^{-1}\psi^{-1} = 1. \end{aligned}$$

Notice that $f_{(\gamma^n, \tau^n)} = \rho_n$ (as defined is Chapter 3) for every $n \in \mathbb{N}$ and so by lemma 3.3, the sequence $f_{(\gamma^n, \tau^n)}$ is eventually faithfull.

The image of every homomorphism $f_{(\gamma^n,\tau^n)}$ contains F, hence the image is dense. Finally, we show that $\chi = \{(\psi,\xi) \in \Omega \mid f_{(\psi,\xi)} \text{ is faithfull}\}$ is residual in the completely metrizable space Ω and in particular, $\chi \neq \varnothing$. For every $g \in \Gamma \smallsetminus \{1\}$ we denote $\chi_g = \{(\psi,\xi) \in \Omega \mid f_{(\psi,\xi)}(g) \neq 1\}$. As in the previous section, χ_g is open as the inverse image of an open set under a continuous map. Since $f_{(\gamma^n,\tau^n)}$ is eventually faithfull there exits $n_0 \in \mathbb{N}$ such that $f_{(\gamma^n,\tau^n)} \in \chi_g$ for all $n \geq n_0$. Since $\langle\tau\rangle$ and $\langle\gamma\rangle$ are non discrete we get from lemma 4.1 that Ω is non-discrete and so by lemma 4.2 we have that $\{(\gamma^n,\tau^n) \mid n \geq n_0\}$ is dense in Ω. This shows that χ_g is dense, hence

$$\chi = \bigcap_{g\in\Gamma\smallsetminus\{1\}} \chi_g$$

is residual.

References

[1] J. Barlev and T. Gelander, *Compactifications and algebraic completions of limit groups*, arXiv:0904.3771v1 (2009).

[2] J. D. Dixon, *Most finitely generated permutation groups are free*, Bull. London Math. Soc. **22** (1990), no. 3, 222–226.

[3] J. Souto E. Breuillard, T. Gelander and P. Storm, *Dense embeddings of surface groups*, Geom. Topol **10** (2006), 1373–1389.

[4] T. P. McDonough, *A permutation representation of a free group*, Quart. J. Math. Oxford Ser. (2) **22** (1977), no. 111, 353–356.